



March 18, 2024

A Word to the Mental Health Consumer from NEFESH **The International Network of Orthodox Mental Health Professionals**

Executive Director
Cynthia R. Darrison, MSILR

President
Rabbi Barry Holzer, MD

Vice President
Karyn Feinberg, PsyD

Secretary
Tzippora Wallach, PsyD

Treasurer
Chana Kahn, LCSW

Past Presidents:
Abe Worenklein, PhD
Nathan A. Solomon, PhD
Mordechai Glick, PhD, z"l
Rabbi Simcha Feuerman, DHL,
LCSW-R

Executive Board:
Ruchama Bistrizky-Clapman
Shalom Feinberg, MD
Rabbi Dovid Fox, PhD
Menachem Hojda LMSW
Rifkie Jungreis, LCSW
Nachum Klafter, MD
Hindie M. Klein, PsyD
Yehuda Krohn, LCSW
Chedva Lax, LCSW
Malkie Schick, LCSW
Chaim Sender, LCSW
Lisa Twerski, LCSW
Rozi Wax, LMFT, LMHC

Rabbinic Advisory Board:
Harav Dovid Cohen

Director of Operations
Alice Tusk, LMSW

In an effort to improve the quality of mental health services to the Torah community, the Board of Directors of Nefesh International has developed a set of guidelines for the consumer. These guidelines are intended to help those seeking mental health services from a **private/independent** practitioner to find a competent provider. On the other hand, psychotherapists in a **clinic** setting should receive clinical supervision from a licensed practitioner.

- Every practitioner should have a minimum of a Master's degree from an accredited school in a clinical mental health field.
- The psychotherapist should be licensed in the state in which services are offered. This license should permit independent clinical practice by the practitioner.
- Masters level clinicians (e.g., LMSW, LMHC, LCSW) in private practice should have completed at least 5 years of 30 hours/week of clinical work. Doctoral level therapists should have at least 2 years post licensure of 30 hours/week of clinical work. Psychiatrists should have completed a residency program accredited by the Accreditation Council for Graduate Medical Education in the field of psychiatry.
- A private clinician in his/her first 10 years of practice should have ongoing supervision from a more experienced colleague.
- The private psychotherapist should have a *rav* of whom s/he asks *sh'eilos*.

The consumer should not be reluctant to ask a potential provider whether he or she meets these standards. It is our hope that these suggestions will encourage better choices in the selection of either **private or clinic-based** practitioners and thus contribute to more successful outcomes. While these guidelines were developed relative to practice in New York and New Jersey, they generally can and should be applied to any geographical area.